

RESEARCH ARTICLE

Diatoms on the carapace of common snapping turtles: *Luticola* spp. dominate despite spatial variation in assemblages

Shelly C. Wu^{1,2*}, Elizabeth A. Bergey^{1,2}

1 Oklahoma Biological Survey, Norman, Oklahoma, United States of America, **2** Department of Biology, University of Oklahoma, Norman, Oklahoma, United States of America

✉ Current address: Andrews Institute, Texas Christian University, Fort Worth, Texas, United States of America

* shelly.wu@tcu.edu



Abstract

Filamentous algae are often visible on the carapaces of freshwater turtles and these algae are dominated by a few species with varying geographic distributions. Compared to filamentous algae, little is known about the much more speciose microalgae on turtles. Our objectives were to compare the diatom flora on a single turtle species (the common snapping turtle, *Chelydra serpentina*) across part of its range to examine spatial patterns and determine whether specific diatom taxa were consistently associated with turtles (as occurs in the filamentous alga *Bacillaria* spp.). Using preserved turtle specimens from museums, we systematically sampled diatoms on the carapaces of 25 snapping turtles across five states. The diverse diatom assemblages formed two groups—the southern Oklahoma group and the northern Illinois/Wisconsin/New York group, with Arkansas not differing from either group. Of the six diatom species found in all five states, four species are widespread, whereas *Luticola* cf. *goeppertiana* and *L.* cf. *mutica* are undescribed species, known only from turtles in our study. *L.* cf. *goeppertiana* comprised 83% of the diatom abundance on Oklahoma turtles and was relatively more abundant on southern turtles (Oklahoma and Arkansas) than on northern turtles (where mean abundance/state was > 10%). *L.* cf. *mutica* was the most abundant species (40%) on New York turtles. Some *Luticola* species are apparently turtle associates and results support a pattern of spatial variation in *Luticola* species, similar to that in *Bacillaria*. Using museum specimens is an efficient and effective method to study the distribution of micro-epibionts.

OPEN ACCESS

Citation: Wu SC, Bergey EA (2017) Diatoms on the carapace of common snapping turtles: *Luticola* spp. dominate despite spatial variation in assemblages. PLoS ONE 12(2): e0171910. doi:10.1371/journal.pone.0171910

Editor: Michael E. Douglas, University of Arkansas Fayetteville, UNITED STATES

Received: December 2, 2016

Accepted: January 28, 2017

Published: February 13, 2017

Copyright: © 2017 Wu, Bergey. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information Files.

Funding: The Collections-Web Award from the National Science Collections Alliance was granted to the corresponding author (SW). The University of Oklahoma Department of Biology and the Oklahoma Biological Survey provided travel support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Introduction

Hard-surfaced, benthic substrates can be a limiting resource for attached organisms in both freshwater and marine aquatic habitats because of the abundance of organisms that can potentially settle and establish on these surfaces [1]. The consequent biofilm varies from thin coatings to upright, architecturally-rich assemblages on rocks and other submerged surfaces [2, 3]. In addition to abiotic substrates, colonized surfaces include aquatic flora and fauna, in which case, the associated biofilm organisms are referred to as epibionts.

Competing interests: The authors have declared that no competing interests exist.

Turtles are an excellent model to study the host-epibiont relationship in freshwaters. Turtles are large and fairly speciose with species varying in habitat use, behavior, and distribution. The most studied alga epibiont on turtles are filamentous green algae in the genus *Basycladia* (= *Arnoldiella* in [4]), species that are rare on other substrates [5, 6]. In the United States, two turtle epibionts (*B. chelonum* and *B. crassa*) often co-occur in the same regions and on the same turtle species [5, 7]. Although the complete distribution of these species is unclear, they are known from Ontario in the north [8] to Cuba in the south [9], and westward to Arizona [10]. Beyond this range, *B. chelonum* has been reported as a non-native species in Oregon [11] and *Basycladia* species, including *B. chelonum*, have been reported on turtles in South America [12, 13]. Other turtle-dwelling *Basycladia* species are found in Japan [14] and Australia [15].

In contrast to macroscopic filamentous algae, microalgae on turtle carapaces have been little studied, but recent reports indicate a combination of generalists and host specialists. Specifically, two new diatom species, *Tursiocola podocnemicola* and *Luticola deniseae* were recently described from turtles in the Amazon Basin in Brazil [16, 17] and a third species, *Mastogloia sterijovskii*, was described from a Macedonian turtle [18]. Floristic surveys of turtle-dwelling diatoms include surveys on the European pond turtle (*Emys orbicularis*) in Turkey [19, 20], with a total of 18 diatom species, a survey of two turtle species (*Pseudemys concinna* and *Trachemys scripta*) in Arkansas (USA), which lists 13 genera [21], and a list of 13 taxa on the red-headed river turtle (*Podocnemis erythrocephala*) in Brazil [17].

In contrast to ecological studies of the turtle-associated *Basycladia* (e.g. [6, 12, 22]), ecological understanding of turtle-associated diatoms is lacking. As a step in understanding the ecology of turtle-associated diatoms, our main objectives were to assess distributional patterns of diatom assemblages across the range on a single turtle species and to determine if there is an association between turtles and any particular diatom species (similar to the association of turtles with *Basycladia*). The common snapping turtle (*Chelydra serpentina*) was chosen to assess diatom assemblages across states because this species hosts (macroscopic) algae on its carapace [23–25] and has a wide distribution, covering two-thirds of the United States [26]. A secondary objective of this project was to trial the use of museum specimens to study turtle epibionts, as our study was based entirely on turtle specimens from two natural history museums.

Materials and methods

Turtle sampling

Regions chosen for this study, based on the availability of museum specimens, were Oklahoma (n = 9 turtles), Arkansas (n = 4), Illinois (n = 5), Wisconsin (n = 4), and New York (n = 3) (Fig 1). Oklahoma turtles were sampled at the Sam Noble Oklahoma Museum of Natural History in Norman, Oklahoma and turtles from other states were sampled at the Field Museum of Natural History in Chicago, Illinois (specimen data is listed S1 Table). Although we had anticipated a larger sample size based on the supplied list of museum specimens, many specimens were unsuitable because they were hatchlings/young juveniles or were road kills with smashed carapaces. Juvenile turtles have relatively few filamentous algae on their carapaces [5, 10], possibly because of their general behavior of hiding during the day, which would limit the light needed for algal growth [27].

Turtle carapaces were sampled systematically. Three vertebral scutes and a total of three costal and marginal scutes (Fig 2) were sampled by placing a plastic tube with a 2.54 cm internal diameter on each sampled scute and algae in the enclosed area were removed by brushing with a test tube brush (diameter: 1.3 cm). The total area sampled on each turtle carapace was 30.4 cm². Samples were preserved in 70% ethanol. Although the use of preserved museum



Fig 1. Origin of snapping turtles used to assess epibiotic algae. Numbers within dots indicate that more one turtle specimen for the location and larger circles with dashed outlines that are centered in states indicate unknown locations within states. The states map shows the majority of the turtle's distribution in North America (e.g., NatureServe 2015). State abbreviations are: OK = Oklahoma; AR = Arkansas; IL = Illinois, WI = Wisconsin, NY = New York.

doi:10.1371/journal.pone.0171910.g001

specimens does not require a permit, the turtle sampling protocol was approved by the University of Oklahoma Institutional Animal Care and Use Committee (tracking number R14-008).

Diatom processing

Samples were processed to eliminate the organic material prior to diatom species identification. Samples were dried onto coverslips and coverslips were heated in a muffle furnace at 450 °C for 1.5 hours. Cooled coverslips were mounted on microscope slides with Naphrax mounting medium (PhycoTech, Inc., St. Joseph, MI). Diatoms were viewed under 1000x magnification using an Olympus CX41 microscope and were identified using [28] and the Diatoms of the United States website [29]. Diatoms were counted to 200 valves by scanning transects across the coverslip. For samples with less than 200 valves, all the diatom valves in the sample were counted.

Statistical analysis

Untransformed diatom abundance and richness data were analyzed with 1-way ANOVA (data met homogeneity and equal variance tests). Tukey tests were used to identify differences in diatom assemblages among states following a significant main test. Raw counts of diatom valves in the assemblage data set were square root transformed prior to multivariate analysis, a step that reduced the weight of less common species [30]. Non-metric Multidimensional Scaling (NMDS) was performed with Bray Curtis similarities. One way-PERMANOVA (Permutational MANOVA) tests, using type III sums of squares and 4999 permutations, were used to test for differences in diatom assemblages among states. For significant PERMANOVA results, associated pair-wise tests were used to identify significantly different states. Similarity Percentage Analysis (SIMPER) was used to identify the diatom taxa contributing the most to differences between states. PERMANOVA, NMDS, and SIMPER analyses were run with PRIMER version 6 and PERMANOVA+ packages (Primer-E Ltd, Plymouth Marine Laboratory, Plymouth, U.K.).

Results

Diatoms on the common snapping turtle across regions

A total of 107 diatom species were found on common snapping turtles across the five sampled states (S2 Table). Overall mean diatom richness was 12.5 ± 1.9 SE taxa per turtle (median = 9;




Fig 2. Areas sampled on the turtle carapace. Green circles represent the sampled locations: three vertebral scutes and three costal/marginal scutes.

doi:10.1371/journal.pone.0171910.g002

range: 2–37 taxa). Only six diatom species (*Caloneis bacillum*, *Gomphonema parvulum*, *Luticola* cf. *goeppertiana*, *Luticola* cf. *mutica*, *Nitzschia amphibia*, and *Nitzschia inconspicua*) were found on snapping turtles in all five states. Of these species, *Luticola* cf. *goeppertiana* occurred on the greatest number of turtles (21 of 25 turtles). In contrast, 59 diatom species (55% of all species found on snapping turtles) occurred in only a single state.

Diatom richness and abundance were variable both among and within states. As a consequence of this within state variation, neither diatom abundance nor richness was significantly different among states (ANOVA: abundance: $F_{4,20} = 2.62$, $P = 0.066$; richness: $F_{4,20} = 1.18$,




Fig 3. Mean (+ SE) abundance (A) and richness (B) values for diatom assemblages on snapping turtles. OK = Oklahoma; AR = Arkansas, IL = Illinois, WI = Wisconsin and NY = New York.

doi:10.1371/journal.pone.0171910.g003

$P = 0.35$, respectively). Trends were evident, however (Fig 3). Arkansas snapping turtles averaging only 20.8 (8.2 SE) diatoms per sample, whereas New York turtles averaged 141.0 (59.0 SE) diatoms. Among states, mean diatom richness per turtle was lower in Arkansas and Oklahoma (7.0 to 7.5 species) than in the other three states (15.6 to 21.3 diatom species per turtle).

Luticola cf. *goeppertiana* was the most common diatom on turtles in three states (OK, AR, IL), where percent abundance ranged from 27% to 83% (Table 1). In Wisconsin, *Frustulia rhomboides* (15%) was the most abundant species and in New York, *Luticola* cf. *mutica* (40%) was the most abundant. Although not the most common species, *Luticola* cf. *goeppertiana* comprised over 10% of the assemblage in both Wisconsin and New York. A few species were slight acidophiles; these included *Tabellaria flocculosa*, 9 species of *Pinnularia*, and 8 species of *Eunotia*. These 18 species were distributed across states as follows: WI (15 spp.) > IL (6 spp.) > NY (4 spp.) > OK (2 spp.) > AR (1 spp.).

Diatom assemblages on the common snapping turtle were significantly different across states (PERMANOVA: pseudo- $F_{4,29} = 2.20$; $P = 0.0002$; Fig 4). Specifically, diatom assemblages on Oklahoma turtles differed from assemblages in Illinois, Wisconsin, and New York (pseudo- t : $P \leq 0.01$), whereas Arkansas diatom assemblages did not differ from Oklahoma or the Illinois-Wisconsin-New York assemblages ($P > 0.05$). Based on SIMPER analysis, Oklahoma turtles had a higher percent abundance of *Luticola* cf. *goeppertiana* (83% of all OK diatoms) compared to Illinois (26.8%), New York (14%) and Wisconsin turtles (11%) (S3 Table; Table 1). Other taxa that contributed to differences between pairs of states include a greater abundance of *Luticola* cf. *mutica* (40%) on New York turtles compared to Oklahoma turtles (1.6%) and the high abundance of *Frustulia rhomboides* (15%) on Wisconsin turtles and this taxon's absence on Oklahoma turtles.

Table 1. Percent composition of numerically dominant diatom taxa sampled on snapping turtles from the five states.

	States				
	OK	AR	IL	WI	NY
<i>Achnanthydium</i> sp. 1			+	7.9	+
<i>Aulacoseira granulata</i>	+		12.8	+	+
<i>Epithemia adnata</i>			7.4	+	
<i>Eunotia</i> sp. 1				5.8	
<i>Frustulia rhomboides</i>				15.2	+
<i>Luticola</i> cf. <i>goeppertiana</i>	83.3	34.9	26.8	10.6	13.9
<i>Luticola</i> cf. <i>mutica</i>	+	+	+	+	40.2
<i>Navicula cryptonella</i>		+	+	5.2	+
<i>Nitzschia frustulum</i>	+	18.1	+		+
<i>Nitzschia inconspicua</i>	+	+	7.4	+	+
<i>Pinnularia microstauron</i>				5.5	
Mean diatom count	68.0	20.8	88.8	82.5	141.0

Percentages are listed for the most common species that cumulatively comprise at least 50% of total diatom abundance on turtles within each state. ‘+’ indicates that the species is present, but not as a numerically dominant in turtles from the other states. Blanks indicate diatom taxa not found on the states’ turtles.

doi:10.1371/journal.pone.0171910.t001

Discussion

Diatom assemblages on snapping turtles from five states showed distinct spatial patterns despite variation in collection dates and years, and differences in collection site location, if known, within states—all of which affect diatom assemblages. Oklahoma diatom assemblages differed from assemblages from the more northern states of Illinois, Wisconsin, and New York, whereas Arkansas assemblages did not differ from either of these groups. The similarity of Oklahoma-Arkansas and Illinois-Wisconsin diatoms could indicate spatial similarity in the diatom floras of adjacent states but the similarity of floras from non-contiguous states (AR versus IL-WI-NY; IL-WI versus NY), indicates other factors than proximity.




Fig 4. Non-metric multi-dimensional scaling plots of diatom assemblages on the shells of snapping turtles collected from five states (Oklahoma, Arkansas, Illinois, Wisconsin, and New York). The plots show the three orthogonal projections (A, B, and C) of the 3-D plot (stress = 0.17).

doi:10.1371/journal.pone.0171910.g004

The general north-south (IL-WI-NY versus OK) difference in turtle diatom flora might be related to latitudinal differences in annual activity period by snapping turtles or biogeographical patterns in diatom distributions. Snapping turtles are occasional aerial baskers [31] that have latitudinal variation in the annual period of activity. For example, Illinois snapping turtles are active for 10 months, from February to December [32], whereas snapping turtles from Ontario, Canada are active for only 4 months, from June to October [33]. This longer activity period among southern turtles may mean greater aerial exposure of carapace algae over the active season. Diatoms are susceptible to desiccation [34] to greater aerial exposure might impact diatom assemblages. Indeed, Oklahoma and Arkansas turtles show a strong trend toward lower diatom abundance and richness than turtles in the three northern states. *Luticola* is considered as an aerophilic to subaerial genus [35–37] and the higher percent abundance of the species *Luticola* cf. *goeppertiana* in OK and AR (35% to 83%, respectively) relative to the northern states (11% to 27%) corresponds to greater aerial exposure over the annual activity period in southern versus northern states. In addition to this latitudinal pattern, diatom biogeography is affected by a combination of spatial and environmental factors [35]. Evidence for biogeographic effects in our study are that 1) 57% of the diatom taxa on snapping turtles were found in only one state and that 2) slightly acidophilic taxa, including *Tabellaria flocculosa*, *Pinnularia* species, and *Eunotia* species, occurred almost entirely in the northern states, indicating effects of regional water chemistry.

Although we found variation in the diatom assemblages among states, a survey of marine olive ridley turtles (*Lepidochelys olivacea*) from an extensive nesting area in Costa Rica found very similar diatom floras on the carapaces [38]. Olive ridley turtles nest periodically over a period of three to five months and may also coexist in feeding areas [39]. Snapping turtles in our study were collected over a wide range of years and sites (including different watersheds and different habitats); consequently, it is not surprising that the turtles in our study displayed greater variation in diatom assemblages than did the marine turtles—including a much greater species richness (our study = 107 diatom species on 25 snapping turtles; [38] = 21 diatom species on 38 olive ridley turtles).

The genus *Luticola* is a characteristic epibiont of snapping turtles. *Luticola* cf. *goeppertiana* occurred on 84% of the common snapping turtles in this study and often reached high abundances. A second species, *Luticola* cf. *mutica* was often present, but only abundant on New York snapping turtles. A third species, *Luticola deniseae*, was described from a Brazilian turtle and as in the *Luticola* in our study, *L. deniseae* dominated the epibiotic diatom flora, comprising over 80% of diatoms. *L. deniseae* was found exclusively on turtles and not on epiphytic and epilithic substrates in the same habitat [17]. One limitation to our study was that we were unable to sample other substrates in collection habitats to determine whether *Luticola* cf. *goeppertiana* occurs selectively on turtles. In contrast to the two encountered *Luticola* species in our study, the other four diatom species found in all five states (*Caloneis bacillum*, *Gomphonema parvulum*, *Nitzschia amphibia*, and *Nitzschia inconspicua*) are widely distributed species occurring on a variety of benthic substrates. More studies are needed to evaluate the degree of specificity of *Luticola*, possibly other diatom taxa on turtles, and other aquatic biota—as well as ecological studies to better understand the nature of the diatom-turtle association.

In contrast to possible specialist species in freshwaters, specialist diatom species have not been apparent on marine turtles. An SEM analysis of diatoms on single specimens of 7 marine turtle species found only 18 diatom taxa, with *Achnanthes* sp. on three turtle species, *Amphora* sp. and *Poulinea* sp. each on two turtle species [40]. Further assessment of the distribution and ecology of epizoic diatoms on both freshwater and marine turtle species may provide insights into the apparent differences in diatom specialization on turtles.

One possibility for the prevalence of *Luticola* on turtles is the ability to tolerate desiccation during aerial exposure when basking. *Luticola* is classed as a subaerial taxon [35] and like many such taxa, has occluded pores in the siliceous cell wall [41], which is an adaptation for reducing water loss [42]. The reduced openings may allow *Luticola* on turtle carapaces to tolerate aerial exposure during basking and terrestrial movement. In an experimental study that tested the tolerance of terrestrial and aquatic diatom taxa (*Luticola* were not tested) to desiccation and high temperature, all tested diatoms succumbed to desiccation but terrestrial diatoms tolerated high temperatures better than aquatic diatoms [34]. Thus, the combination of occluded pores that reduce water loss and probable tolerance to high temperatures during aerial exposure may allow *Luticola* to persist on turtle carapaces exposed to desiccation and high temperatures during basking and terrestrial movements.

Luticola and *Basycladia/Arnoldiella* are both widespread genera that include epibionts on freshwater turtles. Epibiotic species in both genera have varying geographical distributions (though tentative in *Luticola* due to the paucity of studies). Thus far, *Luticola* is known on turtles from only North and South America from this study and [17], respectively, whereas *Basycladia* has a wider distribution, occurring widely in North and South America, and in Australia [15]. Most species of *Basycladia* are turtle epiphytes (one species is found on snails and two typically occur on abiotic substrates: [9, 43]), whereas most species of *Luticola* are not associated with turtles and are more typically found in soil or moss [37].

Our study was greatly facilitated by sampling museum specimens, an approach used previously for epibiotic filamentous algae on turtles [5]. Benefits of using museum collections include a combination of saving time, reducing research costs, avoiding unnecessary duplication of specimens [44], and eliminating stress to live turtles or other organisms. Limitations associated with using museum turtle specimens for studying epizotic diatoms include: 1) exact localities (and associated environmental data) are often unknown; 2) an inability to select random sampling locations or to standardize sampling variables (e.g., date of collection); and 3) possible effects of post-capture turtle processing. For example, post-capture processing would occur if museum personnel removed filamentous algae from the carapaces. Although past history of curation at the Sam Noble Museum of History is not known, algal scrapping is not a current practice and has not been documented in their database (Jessa Watters, personal communication) and algae are not routinely removed from turtles at the Field Museum (Alan Resetar, personal communication). Even if filamentous algae were removed, filamentous algae are often restricted to the sides and posterior of the carapace [5, 11] and sampling six areas of the carapace reduces any impacts on the sampled diatom assemblage. The high species richness of diatoms found in our study using museum specimens (107 species) in comparison to reported diatom richness found on live turtles (e.g. 13 to 18 taxa in [17, 20, 21]) indicates that museum specimens are a good source of epibiotic diatoms. Continuing this approach of using museum turtles and other collected taxa will allow efficient surveys of the poorly known epibionts.

Supporting information

S1 Table. List of turtle specimens sampled to assess diatom assemblages on the carapaces of snapping turtles. Museums are: OMNH = Sam Noble Oklahoma Museum of Natural History, Norman, Oklahoma; FMNH = Field Museum of Natural History, Chicago, Illinois. (DOCX)

S2 Table. Diatom data for snapping turtle samples. (XLSX)

S3 Table. Summary of SIMPER pairwise comparisons of diatom assemblages on shells of snapping turtles. The table shows the five highest-contributing diatoms to differences between pairs of states with significantly different diatom assemblages (OK-IL, OK-WI, and OK-NY). SIMPER analysis was run on square-root transformed data, but data shown are untransformed mean counts (number per sample) for clarity. Diatom species occurring in all states are bolded and values in parentheses are mean counts for these diatoms that were not ranked high in the SIMPER analysis. (DOCX)

Acknowledgments

We thank Cameron Siler, Caryn Vaughn, and Eric Bright for providing advice and comments on versions the manuscript, and Jessa Watters and Alan Resetar for assistance at the Sam Noble Museum and Field Museum, respectively. We also thank two anonymous reviewers for their comments on this manuscript.

Author Contributions

Conceptualization: SW EB.

Formal analysis: SW EB.

Funding acquisition: SW EB.

Investigation: SW.

Methodology: SW EB.

Resources: SW EB.

Visualization: SW EB.

Writing – original draft: SW EB.

References

1. Jackson JBC. Competition on marine hard substrata: the adaptive significance of solitary and colonial strategies. *Am Nat* 1977; 111: 743–767.
2. Golladay SW, Sinsabaugh RL. Biofilm development on leaf and wood surfaces in a boreal river. *Freshwater Biol.* 1991; 25: 437–450.
3. Gorbushina AA. Life on the rocks. *Environ Microbiol.* 2007; 9: 1613–31. doi: [10.1111/j.1462-2920.2007.01301.x](https://doi.org/10.1111/j.1462-2920.2007.01301.x) PMID: [17564597](https://pubmed.ncbi.nlm.nih.gov/17564597/)
4. Boedeker C, O’Kelly CJ, Star W, Leliaert F. Molecular phylogeny and taxonomy of the *Aegagropila* clade (Cladophorales, Ulvophyceae), including the description of *Aegagropilopsis* gen. nov. and *Pseudocladophora* gen. nov.1. *J Phycol.* 2012; 48: 808–825. doi: [10.1111/j.1529-8817.2012.01145.x](https://doi.org/10.1111/j.1529-8817.2012.01145.x) PMID: [27011097](https://pubmed.ncbi.nlm.nih.gov/27011097/)
5. Edgren RA, Edgren MK, Tiffany LH. Some North American turtles and their epizootic algae. *Ecology.* 1953; 34: 733–40.
6. Proctor VW. The growth of *Basycladia* on turtles. *Ecology.* 1958; 39: 634–645.
7. Lipsey LL. *Basycladia chelonum* (Collins) Hoffmann & Tilden and *Basycladia crassa* Hoffmann & Tilden (Cladophorales) in southern Illinois. *Rhodora.* 1978; 80: 441–443.
8. Colt L, Saumure R, Baskinger S. First record of the algal genus *Basycladia* (Chlorophyta, Cladophorales) in Canada. *Can Field Nat.* 1995; 109: 454–455.
9. Normandin RF, Taft CE. A new species of *Basycladia* from the snail *Viviparus malleatus* Reeve. *Ohio J Sci.* 1959; 59: 58–62.
10. Hulse AC. Carapacial and plastral flora and fauna of the Sonora mud turtle, *Kinosternon sonoriense* Le Conte (Reptilia, Testudines, Kinosternidae). *J Herpetol.* 1976; 10: 45–48.

11. Bury RB, Wehr JD, Bury GW, Baggett CL, Doten K. High incidence of filamentous algae on western pond turtles, *Actinemys marmorata*, in the Willamette Valley, Oregon. *Northwest Nat.* 2015; 96: 150–153.
12. Zanelli HRR, Necchi Júnior O, Brites VLC. Spatial and temporal dynamics of *Bacillaria paxillifer* epizoic algae associate with *Phrynops geoffroanus* freshwater turtle in the Uberabinha River, Minas Gerais, Uberlândia. *Arq Ciênc Vet Zool UNIPAR.* 2009; 12: 91–92.
13. Williner V, Carvalho DA, Collins PA. Feeding spectra and activity of the freshwater crab *Trichodactylus kensleyi* (Decapoda: Brachyura: Trichodactylidae) at La Plata basin. *Zoo Stud.* 2014; 53(1): 71.
14. Yoneda Y. Observations on the algae growing on the pond tortoises, with special reference to *Bacillaria crassa* Hoffmann & Tilden. *Publ Seto Mar Biol Lab.* 1952; 2: 227–234.
15. Skinner S, FitzSimmons N, Entwistle TJ. The moss-back alga (Cladophorophyceae, Chlorophyta) on two species of freshwater turtles in the Kimberleys. *Telopea.* 2008; 12: 279–284.
16. Wetzel CE, Van de Vijver B, Cox EJ, Bicudo DdC, Ector L. *Tursiocola podocnemicola* sp. nov., a new epizoic freshwater diatom species from the Rio Negro in the Brazilian Amazon Basin. *Diatom Res.* 2012; 27: 1–8.
17. Wetzel C, Van de Vijver B, Ector L. *Luticola deniseae* sp. nov. a new epizoic diatom from the Rio Negro (Amazon hydrographic basin). *Vie Milieu.* 2010; 60: 177–184.
18. Pavlov A, Jovanovska E, Wetzel CE, Ector L, Levkov Z. Freshwater *Mastogloia* (Bacillariophyceae) taxa from Macedonia, with a description of the epizoic *M. sterijovskii* sp. nov. *Diatom Res.* 2016; 31: 85–112.
19. Soylu E, Gönülol A, Sukatar A, Ayaz D, Tok C. Epizoik freshwater algae on *Emys orbicularis* (Testudinata: Emydidae) from the central Anatolia Region of Turkey. *J Freshwater Ecol.* 2006; 21: 535–538.
20. Ersanli ET, Gonulol A. Epizooiphyte composition on spotted turtle *Emys orbicularis* (Linnaeus, 1758) in Turkey. *Bangladesh J Botany.* 2015; 43: 157–161.
21. Tumlison R, Clark S. Microorganisms associated with the carapace and plastron of aquatic turtles (*Pseudemys concinna* and *Trachemys scripta*) in southwestern Arkansas. *Proc Ark Acad Sci.* 1996; 50: 148–152.
22. Ernst CH, Norris JN. Observations on the algal genus *Bacillaria* and the red-bellied turtle, *Chrysemys rubriventris*. *Estuaries.* 1978; 1: 54–57.
23. Ziglar CL, Anderson RV. Epizoic organisms on turtles in Pool 20 of the upper Mississippi River. *J Freshwater Ecol.* 2005; 20: 389–396.
24. Garbary DJ, Bourque G, Herman TB, McNeil JA. Epizoic algae from freshwater turtles in Nova Scotia. *J Freshwater Ecol.* 2007; 22: 677–685.
25. Belusz LC, Reed RJ. Some epizooiphytes on six turtle species collected in Massachusetts and Michigan. *Am Midl Nat.* 1969; 81: 598–601.
26. Ernst CH, Lovich JE. *Turtles of the United States and Canada*, 2nd ed. Baltimore: John Hopkins University Press; 2009.
27. Neil WT, Allen ER. Algae on turtles: some additional considerations. *Ecology.* 1954; 35: 581–584.
28. Kramer K, Lange-Bertalot H. *Süßwasserflora von Mitteleuropa*. Stuttgart: Gustav Fischer Verlag; 1986–1991.
29. Spaulding SA, Lubinski DJ, Potapova M. *Diatoms of the United States*. 2010. Available from: <http://westerndiatoms.colorado.edu>.
30. Clarke K, Warwick R. *Change in Marine Communities: An approach to statistical analysis and interpretation*. 2nd edition. Plymouth Marine Laboratory, Plymouth, UK: Primer-e Ltd; 2001.
31. Boyer DR. Ecology of the basking habit in turtles. *Ecology.* 1965; 46: 99–118.
32. Smith PW. The amphibians and reptiles of Illinois. *Bull Ill Nat Hist Surv.* 1961; 28: 1–298.
33. Obbard ME, Brooks RJ. A radio-telemetry and mark-recapture study of activity in the common snapping turtle, *Chelydra serpentina*. *Copeia.* 1981; 1981: 630–637.
34. Souffreau C, Vanormelingen P, Verleyen E, Sabbe K, Vyverman W. Tolerance of benthic diatoms from temperate aquatic and terrestrial habitats to experimental desiccation and temperature stress. *Phycologia.* 2010; 49: 309–324.
35. Johansen JR. Diatoms of aerial habitats. In: Smol JP, Stoermer EF, editors. *The diatoms: applications for the environmental and earth sciences*, Second edition. Cambridge, UK: Cambridge University Press; 2010. pp. 465–472.
36. Van de Vijver B, Mataloni G. New and interesting species in the genus *Luticola* D.G. Mann (Bacillariophyta) from Deception Island (South Shetland Islands). *Phycologia.* 2008; 47: 451–467.

37. Spaulding S, Edlund M. *Luticola*. Diatoms of the United States. 2008. Available from: <http://westerndiatoms.colorado.edu/taxa/genus/Luticola>.
38. Majewska R, Santoro M, Bolaños F, Chaves G, De Stefano M. Diatoms and other epibionts associated with olive ridley (*Lepidochelys olivacea*) sea turtles from the Pacific Coast of Costa Rica. *PLoS One*. 2015; 10(6):e0130351. doi: [10.1371/journal.pone.0130351](https://doi.org/10.1371/journal.pone.0130351) PMID: [26083535](https://pubmed.ncbi.nlm.nih.gov/26083535/)
39. Plotkin PT, Byles RA, Rostal DC, Owens DW. Independent versus socially facilitated oceanic migrations of the olive ridley, *Lepidochelys olivacea*. *Marine Biology*. 1995; 122: 137–143.
40. Robinson NJ, Majewska R, Lazo-Wasem EA, Nel R, Paladino FV, Rojas L, et al. Epibiotic diatoms are universally present on all turtle species. *PLoS One*. 2016; 11(6):e0157011. doi: [10.1371/journal.pone.0157011](https://doi.org/10.1371/journal.pone.0157011) PMID: [27257972](https://pubmed.ncbi.nlm.nih.gov/27257972/)
41. Round FE, Crawford RM, Mann DG. *Diatoms: biology and morphology of the genera*. Cambridge, UK: Cambridge University Press; 1990.
42. Lowe RL, Furey PC, Ress JA, Johansen JR. Diatom biodiversity and distribution on wetwalls in Great Smoky Mountains National Park. *Southeast Nat*. 2007; 6:135–152.
43. Peres CK, Branco CCZ. *Basycladia emedii* (Cladophorales, Chlorophyta): A new freshwater epilithic species from Brazil. *Cryptogam, Algal*. 2012; 33: 329–337.
44. Suarez AV, Tsutsui ND. The value of museum collections for research and society. *BioScience*. 2004; 54: 66–74.