A METEORITE FALL IN OKLAHOMA

On the evening of 1936, August 17, at about 7.07 p.m.,

G. S. T., a fireball moved thru the sunlit sky across central
Oklahoma. It was very well viewed in most of Oklahoma and
was also seen from points in Kansas and Texas, and as far south
as waco and Tyler, Texas. A field survey of the fireball was
made immediately by Mr. Robert Brown of the Texas Observers at
Ft. Worth, Texas, and in addition to securing observations which
will be of considerable value in working out the fireball path,
Brown was successful in locating the region of probable fall of
meteorites, and in retrieving one meteorite from the fall four
days later. Two subsequent visits to the region of the endpoint
have been made by Mr. Brown, resulting in fuller information
and in the recovery of one other meteorite from the same fall,
at a point about one mile from where the first one was found.

became conspicuous over the region of Tecumseh, Pottawatomie County, Oklahoma, perhaps a few miles south thereof, at a height of about 26 miles. A few of the better observations seem to refer to an earlier portion of the path. The fireball traveled roughly northwest in azimuth at a low slope (about 11°), and underwent several (perhaps three distinct) explosive flares. Detonations were heard in a region along much of the primarily observed path, in a strip at places almost 100 miles wide. A terminal explosion cloud was formed by the final flare at a height of about 13 miles just northwest of Guthrie, Logan County, and persisted with little drift for 15 minutes or more.

The meteorites were recovered beyond the terminal cloud,

not far from the town of Chescent, for which shey will brokenly be named. The first one was picked up on the third day after the fall, in a fresh condition, by a boy of about 9 named addie Gene Johnson, who was rabbit hunting. Brown was able to lecree and examine the hole this meteorite made on striking, and to check the conditions of its fall. Another party had out fragments off one end of this stone; 1.6 grams of these tiny bits were picked up where they had been allowed to fall when out off, and the major piece as received by us weighed 72.7 grams, and was losf-shaped. The second meteorite was found almost seven weeks after the fall and had gone thru rains; its crust showed a considerably different appearance, being discolored brown by spattered sand and rusting; and cracking of the stone itself had also set in, due to the deteriorating effects of weathering. It weighs only 5.7 grans, and was picked up personally by Mr. Sterling Bunch of the Texas Chservers, who accompanied Brown on third trip of investigation. It is rarely that a meteorite humter has the privilege of picking up his own meteorite .-- but the case demonstrates what large factors interest and knowledge are in the finding of these objects.

possibly of the kind classified as carbonaceous chondrites. They have a low density, not far from 2.0 (less than the density of average terrestrial rocks); the interior ground is generally a dull black, with a few scattered white and gray specks, and some glassy-looking yellowish particles; free metal is relatively sparse, but is well shown by the stone which laid out in several rains, and was detected in the other one by X-ray photographs. The crust of the fresh stone is a shiny black and mostly very thin. The

material is inclined to crumble and cannot be handled freely without some loss.

A fuller description of the fireball is being worked up on behalf of the American Meteor Society, a number of reports having been obtained by letters in addition to those procured thru the field surveys. Later detailed articles will deal with the final fireball data and with proper descriptions and photographs of the meteorites. Further field work is being done to determine more extensive facts about the fall.

Oscaf E. Monnig

and

Robert Brown.

312 W. Loude 3t..

Ft. Worth, Texas.

1936 11-20